arXiv:hep-lat/9908057v1 31 Aug 1999

# Elsevier instructions for the preparation of a 2-column format camera-ready paper in LaTeX

P. de Groot[a]* and X.-Y. Wang[b]

[a]Mathematics and Computer Science Division, Elsevier Science Publishers B.V.,
P.O. Box 103, 1000 AC Amsterdam, The Netherlands

[b]Economics Department, University of Winchester,
2 Finch Road, Winchester, Hampshire P3L T19, United Kingdom

These pages provide you with an example of the layout and style for 100% reproduction which we wish you to adopt during the preparation of your paper. This is the output from the LaTeX document style you requested.

## 1. FORMAT

Text should be produced within the dimensions shown on these pages: each column 7.5 cm wide with 1 cm middle margin, total width of 16 cm and a maximum length of 20.2 cm on first pages and 21 cm on second and following pages. The LaTeX document style uses the maximal stipulated length apart from the following two exceptions (i) LaTeX does not begin a new section directly at the bottom of a page, but transfers the heading to the top of the next page; (ii) LaTeX never (well, hardly ever) exceeds the length of the text area in order to complete a section of text or a paragraph.

### 1.1. Spacing

We normally recommend the use of 1.0 (single) line spacing. However, when typing complicated mathematical text LaTeX automatically increases the space between text lines in order to prevent sub- and superscript fonts overlapping one another and making your printed matter illegible.

### 1.2. Fonts

These instructions have been produced using a 10 point Computer Modern Roman. Other recommended fonts are 10 point Times Roman, New Century Schoolbook, Bookman Light and Palatino.

## 2. PRINTOUT

The most suitable printer is a laser printer. A dot matrix printer should only be used if it possesses an 18 or 24 pin printhead ("letter-quality").

The printout submitted should be an original; a photocopy is not acceptable. Please make use of good quality plain white A4 (or US Letter) paper size. *The dimensions shown here should be strictly adhered to: do not make changes to these dimensions, which are determined by the document style.* The document style leaves at least 3 cm at the top of the page before the head, which contains the page number.

Printers sometimes produce text which contains light and dark streaks, or has considerable lighting variation either between left-hand and right-hand margins or between text heads and bottoms. To achieve optimal reproduction quality, the contrast of text lettering must be uniform, sharp and dark over the whole page and throughout the article.

If corrections are made to the text, print completely new replacement pages. The contrast on these pages should be consistent with the rest of the paper as should text dimensions and font sizes.

---

*Footnotes should appear on the first page only to indicate your present address (if different from your normal address), research grant, sponsoring agency, etc. These are obtained with the `\thanks` command.

Table 1
Biologically treated effluents (mg/l)

| | Pilot plant | | Full scale plant | |
|---|---|---|---|---|
| | Influent | Effluent | Influent | Effluent |
| Total cyanide | 6.5 | 0.35 | 2.0 | 0.30 |
| Method-C cyanide | 4.1 | 0.05 | | 0.02 |
| Thiocyanide | 60.0 | 1.0 | 50.0 | $< 0.10$ |
| Ammonia | 6.0 | 0.50 | | 0.10 |
| Copper | 1.0 | 0.04 | 1.0 | 0.05 |
| Suspended solids | | | | $< 10.0$ |

Reprinted from: G.M. Ritcey, Tailings Management, Elsevier, Amsterdam, 1989, p. 635.

## 3. TABLES AND ILLUSTRATIONS

Tables should be made with LaTeX; illustrations should be originals or sharp prints. They should be arranged throughout the text and preferably be included *on the same page as they are first discussed*. They should have a self-contained caption and be positioned in flush-left alignment with the text margin within the column. If they do not fit into one column they may be placed across both columns (using `\begin{table*}` or `\begin{figure*}` so that they appear at the top of a page).

### 3.1. Tables

Tables should be presented in the form shown in Table 1. Their layout should be consistent throughout.

Horizontal lines should be placed above and below table headings, above the subheadings and at the end of the table above any notes. Vertical lines should be avoided.

If a table is too long to fit onto one page, the table number and headings should be repeated above the continuation of the table. For this you have to reset the table counter with `\addtocounter{table}{-1}`. Alternatively, the table can be turned by 90° ('landscape mode') and spread over two consecutive pages (first an even-numbered, then an odd-numbered one) created by means of `\begin{table}[h]` without a caption. To do this, you prepare the table as a separate LaTeX document and attach the tables to the empty pages with a few spots of suitable glue.

### 3.2. Line drawings

Line drawings should be drawn in India ink on tracing paper with the aid of a stencil or should be glossy prints of the same; computer prepared drawings are also acceptable. They should be attached to your manuscript page, correctly aligned, using suitable glue and *not transparent tape*. When placing a figure at the top of a page, the top of the figure should be at the same level as the bottom of the first text line.

All notations and lettering should be no less than 2 mm high. The use of heavy black, bold lettering should be avoided as this will look unpleasantly dark when printed.

### 3.3. Black and white photographs

Photographs must always be sharp originals (*not screened versions*) and rich in contrast. They will undergo the same reduction as the text and should be pasted on your page in the same way as line drawings.

### 3.4. Colour photographs

Sharp originals (*not transparencies or slides*) should be submitted close to the size expected in publication. Charges for the processing and printing of colour will be passed on to the author(s) of the paper. As costs involved are per page, care should be taken in the selection of size and shape so that two or more illustrations may be fitted together on one page. Please contact the Technical Editor in the Camera-Ready Publications Department at Elsevier for a price quotation and layout instructions before producing your paper in its final form.

Figure 1. Good sharp prints should be used and not (distorted) photocopies.

Figure 2. Remember to keep details clear and large enough.

## 4. EQUATIONS

Equations should be flush-left with the text margin; LaTeX ensures that the equation is preceded and followed by one line of white space. LaTeX provides the document-style option `fleqn` to get the flush-left effect.

$$H_{\alpha\beta}(\omega) = E_{\alpha}^{(0)}(\omega)\delta_{\alpha\beta} + \langle\alpha|W_{\pi}|\beta\rangle \tag{1}$$

You need not put in equation numbers, since this is taken care of automatically. The equation numbers are always consecutive and are printed in parentheses flush with the right-hand margin of the text and level with the last line of the equation. For multi-line equations, use the `eqnarray` environment. For complex mathematics, use the AMS-LaTeX package.

## REFERENCES

1. S. Scholes, Discuss. Faraday Soc. No. 50 (1970) 222.
2. O.V. Mazurin and E.A. Porai-Koshits (eds.), Phase Separation in Glass, North-Holland, Amsterdam, 1984.
3. Y. Dimitriev and E. Kashchieva, J. Mater. Sci. 10 (1975) 1419.
4. D.L. Eaton, Porous Glass Support Material, US Patent No. 3 904 422 (1975).

References should be collected at the end of your paper. Do not begin them on a new page unless this is absolutely necessary. They should be prepared according to the sequential numeric system making sure that all material mentioned is generally available to the reader. Use `\cite` to refer to the entries in the bibliography so that your accumulated list corresponds to the citations made in the text body.

Above we have listed some references according to the sequential numeric system [1–4].

arXiv:hep-lat/9908057v1 31 Aug 1999

# Modelling the Quark Determinant in Full QCD Simulations

A. Duncan[a],E. Eichten[b],and H. Thacker[c]

[a]Dept. of Physics and Astronomy,University of Pittsburgh, Pittsburgh, PA 15260

[b] Fermilab, PO Box 500, Batavia, IL60510

[c]Dept. of Physics, University of Virginia, Charlottesville, VA 22901

The computational requirements and dynamics of Monte Carlo simulations of unquenched QCD incorporating the infrared quark eigenmodes (up to $\approx \Lambda_{QCD}$) exactly and UV modes via a loop representation are discussed. The accuracy of such a loop representation is studied for a variety of lattice volumes and quark masses. The method has been successfully applied for lattices up to $10^3$x20 at $a \simeq$0.17F with improved (clover) action, and allows simulations at or near kappa critical.

## 1. Splitting the Quark Determinant

The essence of the truncated determinant approach [1] to unquenched QCD lies in the realization that the infrared part of the quark determinant (specifically, the determinant $\mathcal{D}(\mathcal{A})$ of the hermitian operator $H \equiv \gamma_5(\not{D}(A) - m))$ can be gauge-invariantly split off, leaving an ultraviolet part which is accurately fit by a linear combination of a small number of Wilson loops. The eigenvalues $\lambda_i$ of $H$

(1) measure quark off-shellness (for $A = 0$, $\lambda_i \rightarrow \pm\sqrt{p^2 + m^2}$),

(2) are gauge-invariant, $\lambda_i(A) = \lambda_i(A^g)$.

Thus we can write $\mathcal{D}(\mathcal{A}) = \mathcal{D}_{\mathcal{IR}}(\mathcal{A})\mathcal{D}_{\mathcal{UV}}(\mathcal{A})$, where the infrared part $\mathcal{D}_{\mathcal{IR}}(\mathcal{A})$ is defined as the product of the lowest $N_\lambda$ positive and negative eigenvalues of $H$, with $|\lambda_i| \leq \Lambda_{QCD}$ (typically, $\simeq$ 300-400 MeV). This cutoff is chosen (a) to include as much as possible of the important low-energy chiral physics of the unquenched theory while (b) leaving the fluctuations of $\ln \mathcal{D}_{\mathcal{IR}}$ of order unity after each sweep updating all links with the pure gauge action. This ensures that the acceptance rate is sufficiently high when the infrared determinant only is used in the accept/reject stage of the procedure. The crucial point is that it is possible to achieve *both* (a) and (b) on fairly large lattices (up to physical volume $\simeq$ 20 $\mathrm{F}^4$) as well as at kappa values arbitrarily close to kappa critical.

## 2. Efficient Computation of the IR spectrum in QCD4

The following Lanczos procedure allows us to extract the needed infrared eigenvalues of $H$ relatively rapidly:

(1) Starting from an initial vector $v_1$, an orthonormal sequence $v_1, v_2, v_3, ..v_{N_L}$ is generated by the standard recursion:

$v_{k+1} = \frac{1}{\beta_k} H v_k - \frac{\beta_{k-1}}{\beta_k} v_{k-1} - \frac{\alpha_k}{\beta_k} v_k$

with the constants $\alpha_k, \beta_k$ determined from overlaps of generated vectors. In the basis of the $v_i$, $H$ is tridiagonal. The corresponding real symmetric tridiagonal matrix $T_{N_L}$ has the $\alpha_k$ on the diagonal and the $\beta_k$ on the sub (and super) diagonal.

(2) A Cullum-Willoughby sieve [2] is used to identify and remove spurious eigenvalues.

(3) The remaining "good" eigenvalues converge most rapidly in the least dense part of the spectrum, in particular, in the needed infrared portion. This remains true even at kappa critical ( in marked contrast to the *matrix inversions* needed in HMC simulations, for example), where the density of eigenvalues near zero is still small. The stability and accuracy of the converged eigenvalues has been checked extensively by gauge transforming the gauge field.

(4) The diagonalization of the $T_{N_L}$ matrix (typically, of order 10,000 in the QCD case) can be completely parallelized using the Sturm sequence property [3] of tridiagonal matrices in

which nonoverlapping parts of the spectrum are independently extracted by a bisection procedure.

## 3. Fitting the UV modes- Loop representations of the Quark determinant

The effective gauge action generated by internal quark loops is a gauge-invariant functional, which can be evaluated explicitly in a hopping parameter expansion:

$$\begin{aligned}\ln\mathcal{D}(A)/V &= 288\kappa^4\sum L_1 + 2304\kappa^6\sum L_2 \\ &+ 4608\kappa^6\sum L_3 + 1536\kappa^6\sum L_4 + ..\end{aligned}$$

where $V$ is the lattice volume, $L_1$ a generic plaquette, and $L_{2,3,4}$ are 6 link loops with link directions $(i,i,j,-i,-i,-j)$, $(i,j,k,-j,-i,-k)$, and $(i,j,k,-i,-j,-k)$ $(1 \leq i,j,k \leq 4)$. Unfortunately, this hopping parameter expansion is useless [4] except for extremely heavy quarks ($\kappa \to 0$), due to the contribution of large loops. Instead, large loops may be cut off by removing the IR modes $|\lambda_i| < \Lambda_{QCD}$. Now the expansion converges much more quickly, and we may write $\ln\mathcal{D}_{UV}(A) = V\sum L_i(A)$ where the sum involves only a small number of loops (as we shall see, typically less than 10) and the coefficients $c_i$ are determined nonperturbatively. Of course, to compute $\mathcal{D}_{UV}$, we need the complete spectra for an ensemble of configurations. The computational cost for extracting a complete Dirac spectrum via Lanczos is large but manageable, as this calculation need only be done for a limited number of decorrelated configurations. For example, the complete Dirac spectrum for a $10^3$x20 lattice has 240,000 eigenvalues and requires about 500,000 Lanczos sweeps, equivalent to about 1 400-Mhz-Pentium-week. The final spectrum can be checked with analytic spectral sum rules which give the sum of powers of the eigenvalues as explicit functionals of small loops.

The results of a fit of $\ln\mathcal{D}_{UV}$ to a linear combination of Wilson loops on an ensemble of $10^3$x20 lattices at beta=5.7 (with clover improvement) and $\kappa$=0.1425 is shown in Fig.1 The configurations were generated including the truncated determinant $\ln\mathcal{D}_{IR}$ and therefore already contain the exact low-energy chiral physics. The fit is very good once 4, 6 and 3 figure-8 8-link operators are included in the fit . These results suggest that the full determinant can be accurately modelled by computing the low eigenvalues exactly and including the remaining high modes via an approximate loop action. In any event, we expect that the UV fluctuations affect primarily the scale of the theory, while for the low energy spectrum, quark off-shellness is limited to about $\Lambda_{QCD}$ and dimensionless mass ratios should therefore be largely insensitive to $\mathcal{D}_{UV}$.



Figure 1. Fit of UV Determinant to Loop Action, $\kappa$=0.1425

## 4. MonteCarlo Dynamics for QCD4 Simulations with the Truncated Determinant

In the truncated determinant approach to QCD4 [1], unquenched configurations are generated by the following algorithm:

1. Update the gauge configuration with the pure gauge action, using a procedure compliant with detailed balance. We have used Metropolis link updates applied to a randomly chosen block of noninterfering links to ensure detailed balance while maintaining parallelizability of the computation.

2. Apply a metropolis accept/reject criterion based on the the effective quark action

$$S_{\text{quark}} = N_F \ln\det\mathcal{D}_{IR}(A) \quad (N_F = 2) \qquad (1)$$



Figure 2. Determinant relaxation, $\kappa$=0.1440

Typically, we use an IR cutoff for the truncated determinant corresponding to a gauge-invariant eigenvalue of 300-400 MeV. This procedure leads to tolerable acceptance rates on lattices of fairly large physical volume - we have explored systems up to 20 $F^4$, while controllable finite size errors in electromagnetic fine structure lattice studies [5] with a long range massless U(1) field require lattice volumes $\geq$6 $F^4$. The simulations in progress are on three different lattice sizes:

(1) $10^3$x20 lattices at $\beta$=5.7 (clover improved) at $\kappa$=0.1415, 0.1425, 0.1436 and 0.1440 (the last value being very close to kappa critical). This corresponds to a physical volume of roughly $(1.7\mathrm{F})^3$x3.4F.

(2) $6^4$ lattices at an effective $\beta$ of 4.5, but using the O($a^2$) improved gauge action of Alford et al,[6], at kappa critical.

(3) $8^4$ lattices at an effective $\beta$=4.5 (O($a^2$) improved), again at kappa critical.

For the $10^3$x20 lattices we have checked that the truncated determinant simulations succeed in equilibrating the configurations, and that they decorrelate reasonably rapidly subsequent to equilibration. The equilibration can be studied by looking at the relaxation of $S_{\mathrm{quark}}$ from the quenched value corresponding to the starting configuration. Even at $\kappa = \kappa_c$, the $10^3$x20 lattices equilibrate after a few hundred sweeps (see Fig. 2).

To measure decorrelation we have calculated the autocorrelation of the pion propagator at various time separations, keeping configurations separated by 20 steps of the basic algorithm. A typical example, from the $10^3$x20 runs with $\kappa$=0.1425, is shown in Fig.3, where the pion correlator at time slice 6 is seen to be effectively decorrelated after about 30 steps of the algorithm.



Figure 3. Decorrelation of pion propagator, $\kappa$=0.1425

Cases (2,3) represent physically large lattices (33 $F^4$ and 105 $F^4$ resp.) at kappa critical and display critical slowing down (several thousand sweeps are needed to equilibrate the configurations). However the lattice sizes are small: new configurations can be generated and the truncated determinant computed quite rapidly ($\simeq$ 20 minutes for the $6^4$ lattices, two hours for the $8^4$ lattices, on a Pentium 400 Mhz processor). These lattices will allow a detailed study of string-breaking and other unquenched dynamical effects (see talk of Eichten, this conference).

## REFERENCES

1. A. Duncan, E. Eichten and H. Thacker, Phys. Rev. D59, 014505 (1998).
2. J. Cullum and R.A. Willoughby, J. Comp. Phys. 44, 329 (1981).
3. G.H. Golub and C.F. Loan, *Matrix Computations*, 2nd edition (Johns Hopkins, 1990).
4. A. Duncan, E. Eichten, R. Roskies and H. Thacker, Phys. Rev. D60, 054505(1999).
5. A. Duncan, E. Eichten, and H. Thacker, Phys. Rev. Lett. 76, 3894 (1996).
6. M. Alford, W. Dimm, G.P. Lepage, G. Hockney, and P.B. Mackenzie, Phys. Lett. B361, 87(1995).